

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Maxillofacial Imaging

Journal homepage: <https://www.ijmi.in/>

Mini Review

Role of physical therapy in temporomandibular joint pain— A short review

VV Manjula Kumari¹, Mohammed Sheeba Kauser^{2*}, Himanshu D Tiwari^{1,3},
Sayed Danial Seyed Mazhari⁴

¹Varanaa's Health care Research and Training Organization LLP, Nellore, Andhra Pradesh, India

²AR ISHAQ Physiotherapy Clinics, Nellore, Andhra Pradesh, India

³69 Fitness Street

⁴Gulf Medical University, UAE



ARTICLE INFO

Article history:

Received 03-01-2024

Accepted 21-02-2024

Available online 14-05-2024

Keywords:

Temporomandibular joint

Pain

Physical therapy

ABSTRACT

This efficient survey broke down investigations looking at the adequacy of different active recuperation intercessions for temporomandibular joint (TMJ) pain. The accompanying suggestions emerged from the various surveys: (1) dynamic activities and manual activations may be compelling; (2) postural preparation might be utilized in blend with different intercessions, as autonomous impacts of postural preparation are obscure; (3) mid-laser treatment might be more compelling than other electrotherapy modalities; (4) programs including unwinding strategies and biofeedback, electromyography preparing, and proprioceptive re-schooling might be more successful than fake treatment or occlusal supports; and (5) blends of dynamic activities, manual treatment, postural adjustment, and unwinding procedures might be viable. These studies ought to be seen carefully.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

TMJ and the masticatory muscle is present almost in 20% of the populace is impacted, with 10% to 20% of those looking for treatment.¹⁻³ These issues moreover are alluded to as "temporomandibular disorder," "craniomandibular messes," and "mandibular dysfunction."^{4,5} The introducing side effects of TMD are: (1) Discontinuous or on the other hand determined torment in the masticatory muscles or the TMJ, furthermore, less often in nearby designs; (2) limits or then again deviations of mandibular development; (3) TMJ sounds six different side effects, like tinnitus, strange gulping, and hyoid bone delicacy, moreover may occur. Personal satisfaction might be impacted, with a pessimistic impact on friendly capability, profound wellbeing, and energy level.⁶⁻⁹

* Corresponding author.

E-mail address: sheebaishaq.doc@gmail.com (M. S. Kauser).

Presently, there is absence of agreement among analysts as to aetiology, analysis, and the executives of this problem. The conclusion of TMD is normally based on the introducing signs and symptoms. The exploration demonstrative rules for temporomandibular problems (RDC/TMD) applies a double hub framework to analyse and arrange patients with TMD. The principal pivot is separated into three groups of usually happening TMDs:

1. Muscle problems, incorporating myofascial treatment with or without restricted mandibular opening.
2. Plate relocation regardless of decrease or restricted mandibular opening.
3. Arthralgia, joint pain, and arthrosis.

A wide assortment of exercise-based recuperation strategies, counting joint preparation, practice remedy, electrotherapy, instruction, biofeedback and unwinding, and

postural revision, have been utilized in the administration of this disorder. Research assessing the impacts of active recuperation in the executives of TMD has been scrutinized for its absence of systemic rigor.^{10–15} Notwithstanding, late investigations have endeavored to address a few recently recognized limits. Since a significant part of the examination inspecting the impacts of active recuperation on TMD has not been distributed in non-intrusive treatment diaries, fostering a proof base for overseeing TMD is difficult.

This deliberate audit of randomized controlled preliminaries (RCTs) and non-randomized controlled preliminaries surveyed the non-intrusive treatment the executives of intense and ongoing TMD on clinically important results like torment, range of movement (ROM), inability and capability, joint commotion, delicacy, and mental elements.^{16–18}

2. Method

Studies from 1980 through January 2011 were taken. Record Medicus (MEDLINE), PUBMED, United Wellbeing Writing (CINAHL), Research Gate, and the Cochrane Focal Register of Controlled Preliminaries were looked through utilizing the text words "facial pain," "physical treatment," "recovery," "temporomandibular jumble (TMD)," "temporomandibular joint (TMJ)," "temporomandibular joint disorder," TMJ dysfunction and "treatment."

3. Discussion

All the authors equally studied the articles thoroughly and found only 30 studies related to our topic. Rest of the articles were set aside. Studies shown during treatment with most types of exercise based recuperation, counting fake treatment. Non-intrusive treatment was accounted for as quite often better than no treatment, with viability expanding in direct extent to how much treatment got. Also, those subjects who got more treatment modalities appeared to show improvement over the people who got fewer modalities.^{19,20} A 1996 deliberate review expressed that there was inadequate proof to disprove or uphold either control or activation in treatment of the TMJ. A later deliberate survey of low-level laser therapy showed a decrease in pain and improvement in well-being status in constant joint problems.^{21–23}

In any case, an efficient survey of ultrasound in the administration of ongoing outer muscle disorders showed little proof to help its utilization. A meta-analysis reasoned that, albeit restricted in degree, the accessible information support the viability of EMG biofeedback medicines for TMD. Incorporation rules differed among the examinations we audited, possible because of the absence of agreement with respect to the conclusion of TMD. The absence

of normalized consideration models is a restriction while contrasting investigations, as well as with regard to the suggestions made. Subjects with myofascial TMD were remembered for 65% of the examinations chosen. Most of the patients who looked for treatment for TMD and were hence engaged with the investigations were women.²⁴ This finding might connect with a distinction in treatment-chasing conduct among people, as well as the more prominent probability for ladies to have somatization disorders. The outside legitimacy of the suggestions is restricted, due to a limited extent, to the distinctions in the gatherings considered. One study showed that the patients who wouldn't take treatment had more agony also, more condition-related impedance in day-to-day existence.

Ladies and men who create persistent TMD show more psychosocial trouble than those whose intense TMD settle. Different indicators of chronicity are TMD of the myofascial type and being female.^{25–27} The utilization of various mediations in a number of studies brought about suggestions based on a multi-mediation program in light of the fact that the viability of a solitary mediation alone was not inspected. A range of various result measures was utilized in the investigations audited. The greater part of the examinations included somewhere in the range of 2 to 5 result measures. In spite of the fact that there was some congruity in the result regions evaluated, the real measures varied among the examinations, with more than 60 various techniques used to survey the results.

Most of the remaining examinations neglected to report either dependability or legitimacy for the result measures utilized, making less trust in the review results. The significance of long haul follow-up to survey the maintenance of present moment treatment impacts is basic to looking at the viability of the intercessions in question.

4. Conclusion

Programs including unwinding procedures and biofeedback, EMG preparing, proprioceptive correction might be more powerful than fake treatment or occlusal braces in diminishing agony and expanding in individuals with intense or persistent myofascial or solid TMD for the time being and the long haul.^{28–30}

Projects including blends of dynamic activities, manual treatment, postural revision, and unwinding procedures might diminish torment and impedance and increment in the present moment in individuals with TMD coming about because of intense plate dislodging, intense joint pain, or intense myofascial TMD.^{31–35}

Be that as it may, it is difficult to perceive whether a mix program is more compelling than giving the different components of the program as individual treatment methods.

5. Source of Funding

None

6. Conflict of Interest


None.

References

- Fabio RD. Physical therapy for patient with TMD: a descriptive study of treatment, disability, and health status. *J Orofac Pain*. 1998;12(2):124–35.
- Pedroni CR, De Oliveira A, Guaratini MI. Prevalence study of signs and symptoms of temporomandibular disorders in university students. *J Oral Rehabil*. 2003;30(3):283–9.
- Nassif NJ, Al-Salleh F, Al-Admawi M. The prevalence and treatment needs of symptoms and signs of temporomandibular disorders among young adult males. *J Oral Rehabil*. 2003;30(9):944–50.
- Taylor M, Suvinen T, Reade P. The effect of Grade IV distraction mobilization on patients with temporomandibular pain-dysfunction disorder. *Physiother Theor Pract*. 1994;10(3):129–36.
- Bertolucci LE, Gray T. Clinical analysis of mid-laser versus placebo treatment of arthralgic TMJ degenerative joints. *Cranio*. 1995;13(1):26–9.
- Dworkin SF, Huggins KH, Wilson L, Mancl L, Turner J, Massoth D, et al. A randomized clinical trial using research diagnostic criteria for temporomandibular disorders-axis II to target clinic cases for a tailored self-care TMD treatment program. *J Orofac Pain*. 2002;16(1):48–63.
- Hall LJ. Physical therapy results for 178 patients with temporomandibular joint syndrome. *Am J Otol*. 1984;5(3):183–96.
- Ali HM. Diagnostic criteria for temporomandibular joint disorders: a physiotherapist's perspective. *Physiother*. 2002;88(7):421–6.
- Carmeli E, Sheklow SL, Blommenfeld I. Comparative study of repositioning splint therapy and passive manual range of motion techniques for anterior displaced temporomandibular discs with unstable excursive reduction. *Physiotherapy*. 2001;87(1):26–36.
- Dworkin SF, Leresche L. Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. *J Craniomandib Disord*. 1992;6(4):301–55.
- Syrop SB. Initial management of temporomandibular disorders. *Dent Today*. 2002;21(8):52–7.
- Glass EG, Glaros AG, Msglynn FD. Myofascial pain dysfunction: treatments used by ADA members. *Cranio*. 1993;11(1):25–9.
- Sturdivant J, Friction JR. Physical therapy for temporomandibular disorders and orofacial pain. *Curr Opin Dent*. 1991;1(4):485–96.
- Feine JS, Widmer CG, Lund JP. Physical therapy: a critique. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 1997;83(1):123–7.
- Feine JS, Lund JP. An assessment of the efficacy of physical therapy and physical modalities for the control of chronic musculoskeletal pain. *Pain*. 1997;71(1):5–23.
- Barry JM. Evidence-based practice in pediatric physical therapy. *PT Magazine*. 2001;9(11):38–51.
- Sackett DL. Rules of evidence and clinical recommendations for the use of antithrombotic agents. *Chest*. 1989;95(2):2–4.
- Megens A, Harris SR. Physical therapist management of lymphedema following treatment for breast cancer: a critical review of its effectiveness. *Phys Ther*. 1998;78(12):1302–11.
- Harris SR. How should treatments be critiqued for scientific merit? *Phys Ther*. 1996;76(2):175–81.
- Law M, Stewart D, Pollock N. Guidelines for critical review form-quantitative studies; 1998. Available from: <http://www.fhs.mcmaster.ca/rehab/ebp/AccessedFebruary20,2003..>
- Burgess JA, Sommers EE, Truelove EL, Dworkin SF. Short-term effects of two therapeutic methods on myofascial pain and dysfunction of the masticatory system. *J Prosthet Dent*. 1998;60(5):606–10.
- De Laat A, Stappaerts K, Papy S. Counseling and physical therapy as treatment for myofascial pain of the masticatory system. *J Orofac Pain*. 2003;17(1):42–9.
- Jagger RG. Mandibular manipulation of anterior disc displacement without reduction. *J Oral Rehabil*. 1991;18(6):497–500.
- Magnusson T, Syren M. Therapeutic jaw exercises and interocclusal appliance therapy. *Swed Dent*. 1999;23(1):27–37.
- Michelotti A, Steenks MH, Farella M, Parisini F, Cimino R, Martina R. The additional value of a home physical therapy regimen versus patient education only for the short-term treatment of myofascial pain of the jaw muscles: short-term results of a randomized clinical trial. *J Orofac Pain*. 2004;18(2):114–25.
- Minagi S, Nozaki S, Sato T, Tsuru H. A manipulation technique for treatment of anterior disk displacement with reduction. *J Prosthet Dent*. 1991;65(5):686–91.
- Nicolakis P, Erdogmus CB, Koff A, Nicolakis M, Piehslinger E, Fialka-Moser V. Effectiveness of exercise therapy in patients with myofascial pain dysfunction syndrome. *J Oral Rehabil*. 2002;29(4):362–8.
- Nicolakis P, Erdogmus CB, Kollmitzer J. An investigation of the effectiveness of exercise and manual therapy in treating symptoms of TMJ osteoarthritis. *Cranio*. 2001;19(1):26–32.
- Nicolakis P, Erdogmus CB, Kollmitzer J. Long-term outcome after treatment of temporomandibular joint osteoarthritis with exercise and manual therapy. *Cranio*. 2002;20(1):23–27.
- Nicolakis P, Erdogmus C, Kopf A, Ebenbichler G, Kollmitzer J, Piehslinger E, et al. Effectiveness of exercise therapy in patients with internal derangement of the temporomandibular joint. *J Oral Rehabil*. 2001;28(12):1158–64.
- Nicolakis P, Erdogmus B, Kopf A, Djaber-Ansari A, Piehslinger E, Fialka-Moser V. Exercise therapy for craniomandibular disorders. *Arch Phys Med Rehabil*. 2000;81(9):1137–42.
- Tegelberg A, Kopp S. Short-term effect of physical training on temporomandibular joint disorder in individuals with rheumatoid arthritis and ankylosing spondylitis. *Acta Odontol Scand*. 1988;46(1):49–56.
- Tegelberg A, Kopp S. A 3-year follow-up of temporomandibular disorders in rheumatoid arthritis and ankylosing spondylitis. *Acta Odontol Scand*. 1996;54(1):14–8.
- Wright EF, Domenech MA, Jr JF. Usefulness of posture training for patients with temporomandibular disorders. *J Am Dent Assoc*. 2000;131(2):202–10.
- Al-Badawi EA, Mehta N, Forgione AG, Lobo SL, Zawawi KH. Efficacy of pulsed radio frequency energy therapy in temporomandibular joint pain and dysfunction. *Cranio*. 2004;22(1):10–20.

Author biography

VV Manjula Kumari, CEO and Senior Consultant Physiotherapist for Obesity

Mohammed Sheeba Kauser, Chief Physiotherapist, Founder & Director  <https://orcid.org/0000-0001-5301-3559>

Himanshu D Tiwari, CEO & Director

Sayed Danial Seyed Mazhari, DMD

Cite this article: Manjula Kumari VV, Kauser MS, Tiwari HD, Mazhari SDS. Role of physical therapy in temporomandibular joint pain— A short review. *IP Int J Maxillofac Imaging* 2024;10(1):11-13.