

An Unusually Large Submandibular Salivary Stone: A Case Report

Shishir Mohan¹, Hasti Kankariya², Arti Chauhan³, Mohit Aggarwal⁴, Akhilanand Chaurasia⁵

^{1,2,3,4}Department of Oral & Maxillofacial Surgery, K.D. Dental College & Hospital, Mathura, ⁵Faculty of Dental Sciences, King George Medical University, Lucknow

***Corresponding Author:**

Email:

Abstract

Sialolithiasis accounts for the most etiology of salivary gland obstruction which leads to recurrent painful swelling of the involved gland which often exacerbates while eating. Stones may be encountered in any of the salivary glands but most frequently in the submandibular gland and its duct. Presented here is a case report of a twenty five year old male patient who had a submandibular sialolith. The sialolith was removed with intraoral approach and no postoperative complications were noted. The article also reviews the various available diagnostic modalities and treatment options.

Keywords: Sialolith, Swelling, Submandibular gland, Laser

Introduction

Sialolithiasis is the most common disease of salivary glands. It is estimated that it affects 12 in 1000 of the adult population¹. Males are affected twice as much as females². Children are rarely affected but a review of the literature reveals 100 cases of submandibular calculi in children aged 3 weeks to 15 years old³. Sialolithiasis accounts for more than 50% of diseases of the large salivary glands and is thus the most common cause of acute and chronic infections⁴. More than 80% occur in the submandibular gland or its duct, 6% in the parotid gland and 2% in the sublingual gland or minor salivary glands. Multiple calculi in the sub-mandibular gland are rare⁵, as is simultaneous lithiasis in more than one salivary gland. 40% of parotid and 20% of submandibular stones are not radiopaque and sialography may be required to locate them². Salivary calculi are usually unilateral and are not a cause of dry mouth. Clinically they are round or ovoid, rough or smooth and of a yellowish colour. They consist of with smaller amounts of magnesium, potassium and ammonia. This mix is distributed evenly throughout⁶. Submandibular stones are 82% inorganic and 18% organic material whereas parotid stones are composed of 49% inorganic and 51% organic material. The organic material is composed of various carbohydrates and amino

acids⁶. Bacterial elements have not been identified at the core of a sialolith⁶.

Case Report

A 25-year-old male presented at the department of Oral & Maxillofacial surgery of K D Dental College & Hospital following referral by general dental practitioner for an opinion on a firm mass in the anterior part of the right side of the floor of the mouth. Patient complain of increase pain during chewing. Extra-oral examination revealed a palpable left submandibular gland and intraoral examination revealed a large, firm, non-tender swelling in the left anterior floor of mouth in the region of the submandibular duct. A lower occlusal radiograph showed the mass to be radiopaque and extending back beyond the lower right first permanent molar (Fig. 1). A diagnosis of left submandibular duct calculus was made and at a subsequent appointment the stone was removed under local anaesthetic with sharp dissection (Fig. 2). It was measured to be 30mm long along its greatest length (Fig. 3). Due to the swelling in the left submandibular triangle and the size of the stone, the patient was reviewed two weeks post operatively to check salivary function of the gland. On review occlusal x-ray was taken (Fig. 4) the left submandibular gland was palpable but clear saliva could be expressed from the duct on massage.



Fig. 1: Mandibular Occlusal x ray shows left submandibular duct stone extending well beyond 1st molar



Fig. 4: Post-operative mandibular occlusal x ray



Fig. 2: Dissection done & stone identified




Fig. 3: Calculi length more than 35mm in length

Discussion

Although large sialoliths have been reported in the body of salivary glands⁷⁻¹⁰, they have been rarely been reported in the salivary ducts^{11,12-14}. Messerly removed a 51 mm long calculus that occupied the entire length of Stenson's duct in a 66-year-old man¹³. Brusati and Fiamminghi removed a sialolith from the left submandibular duct of a 55 year-old man measuring 27x31 mm¹⁴. More recently Leung et al. removed a sialolith 14x9 mm from the right submandibular duct¹. The sialolith removed in our case was comparable to these.

The exact aetiology and pathogenesis of salivary calculi is largely unknown. Genesis of calculi lies in the relative stagnation of calcium rich saliva. They are thought to occur as a result of deposition of calcium salts around an initial organic nidus consisting of altered salivary mucins, bacteria and desquamated epithelial cells¹⁵. For stone formation it is likely that intermittent stasis produces a change in the mucoid element of saliva, which forms a gel. This gel produces the framework for deposition of salts and organic substances creating a stone.

Traditional theories suggest that the formation occurs in two phases: a central core and a layered periphery¹⁶. The central core is formed by the precipitation of salts, which are bound by certain organic substances. The second phase consists of the layered deposition of organic and non organic material¹⁷. Submandibular stones are thought to form around a nidus of mucous¹⁸, whereas parotid stones are thought to form most often around a nidus of inflammatory cells or a foreign body^{18,19,20}. Another theory has proposed that an

unknown metabolic phenomenon can increase the saliva bicarbonate content, which alters calcium phosphate solubility and leads to precipitation of calcium and phosphate ions²¹. A retrograde theory for sialolithiasis has also been proposed¹⁶. Aliments, substances or bacteria within the oral cavity might migrate into the salivary ducts and become the nidus for further calcification. Salivary stagnation, increased alkalinity of saliva, infection or inflammation of the salivary duct or gland, and physical trauma to salivary duct or gland may predispose to calculus formation. Submandibular sialolithiasis is more common as its saliva is (i) more alkaline, (ii) has an increased concentration of calcium and phosphate, and (iii) has a higher mucous content than saliva of the parotid and sublingual glands. In addition, the submandibular duct is longer and the gland has an antigravity flow. Stone formation is not associated with systemic abnormalities of calcium metabolism. Electrolytes and parathyroid hormone studies in patients with sialolithiasis have not shown abnormalities²³. Gout is the only systemic illness known to predispose to salivary stone formation²³, although in gout the stones are made predominantly of uric acid⁶. The proposed association between hard water areas and salivary calculi has been shown to be incorrect²⁴. The lack of association holds equally for both sexes²⁴. One study has suggested a link between sialolithiasis and nephrolithiasis, reporting an association in up to 10% of patients²⁵.

Calculi may cause stasis of saliva, leading to bacterial ascent into the parenchyma of the gland, and therefore infection, pain and swelling of the gland. Some may be asymptomatic until the stone passes forward and can be palpated in the duct or seen at the duct orifice. It may be possible that obstruction caused by large calculi is sometimes asymptomatic as obstruction is not complete and some saliva manages to seep through or around the calculus. Long term obstruction in the absence of infection can lead to atrophy of the gland with resultant lack of secretory function and ultimately fibrosis. Complete obstruction causes constant pain and swelling, pus may be seen draining from the duct and signs of systemic infection may be present.

Bimanual palpation of the gland itself can be useful, as a uniformly firm and hard gland suggests a hypo-functional or non-functional gland. For parotid stones, careful intraoral palpation around Stenson's duct orifice may reveal

a stone. Deeper parotid stones are often not palpable. When minor salivary glands are involved they are usually in the buccal mucosa or upper lip, forming a firm nodule that may mimic tumour²⁶.

It is very uncommon for patients to have a combination of radiopaque and radiolucent stones²⁷; 40% of parotid stones may be radiolucent. Sialography is thus useful in patients showing signs of sialadenitis related to radiolucent stones or deep submandibular/parotid stones. Sialography is, however, contraindicated in acute infection or in significant patient contrast allergy⁶. Treatment Patients presenting with sialolithiasis may benefit from a trial of conservative management, especially if the stone is small.

The patient must be well hydrated and the can apply moist warm heat and gland massage, while sialogogues are used to promote saliva production and flush the stone out of the duct. With gland swelling and sialolithiasis, infection should be assumed and a penicillinase resistant anti-staphylococcal antibiotic prescribed. Most stones will respond to such a regimen, combined with simple sialolithotomy when required^{18,19}. Almost half of the submandibular calculi in the distal third of the duct and are amenable to simple surgical release through an incision in the floor of the mouth, which is relatively simple to perform and not usually associated with complications²⁸. If the stone is sufficiently forward it can be milked and manipulated through the duct orifice. This can be done with the aid of lacrimal probes and dilators to open the duct. Once open, the stone can be identified, milked forward, grasped and removed. The gland is then milked to remove any other debris in the more posterior portion of the duct.

The duct may need opening to retrieve the stone. This involves a trans-oral approach where an incision is made directly onto the stone. In this way more posterior stones, 1–2 cm from the punctum, can be removed by cutting directly onto the stone in the longitudinal axis of the duct. Care is taken as the lingual nerve lies deep, but in close association with the submandibular duct posteriorly. Subsequently, the stone can be grasped and removed. No closure is done leaving the duct open for drainage. If the gland has been damaged by recurrent infection and fibrosis, or calculi have formed within the gland, it may require removal. Parotid stone management is more problematic as only a small segment of Stenson's duct is approachable through an intraoral incision. In addition, opening Stenson's

duct can be complicated by subsequent stenosis of the duct whereas this is rare in the submandibular gland. As a result, parotidectomy is the main stay of surgical management for the majority of intra glandular stones. This is reserved for patients whose symptoms do not respond to conservative therapy and suffer from recurrent pain and swelling. Alternative methods of treatment have emerged such as the use of extracorporeal shock wave lithotripsy (ESWL) and more recently the use of endoscopic intracorporeal shockwave lithotripsy (EISWL), in which shockwaves are delivered directly to the surface of the stone lodged within the duct without damaging adjacent tissue (piezoelectric principle)²⁹. Both extra and intracorporeal lithotripsy are gaining increasing importance in the treatment of salivary stone disease⁴. In extracorporeal piezoelectric lithotripsy, the average size of fragments produced is about 0.7 mm²². Duct diameters are greater than 0.7 mm in general except for at the ostium. Therefore, fragments produced by ESWL would not be prohibited by duct diameters. Findings have also suggested that best results in salivary stone lithotripsy are achieved when the maximum size of stone fragments does not exceed 1.2 mm²². Extracorporeal salivary lithotripsy provides another therapeutic option that carries fewer risks than surgical removal of the affected gland, such as the risks of a general anaesthetic, facial nerve damage, surgical scar, Frey's syndrome, and causes little discomfort to the patient whilst preserving the gland²⁹. A retrospective study of patients treated endoscopically from 1994 to 1999 showed a success rate of 83% with no severe complications³⁰. Endoscopy is a minimally invasive technique for removal of calculi from salivary glands as well as an excellent diagnostic procedure³¹, as miniaturized endoscopes conforming to the physiological widths of the ducts are used to directly view and then deliver shock waves to the stones.

Conclusion

There are various methods available for the management of salivary stones, depending on the gland affected and stone location. These have been mentioned in the preceding paragraphs. It must be noted, however, that ESWL and EISWL offer alternatives to gland removal. Submandibular gland removal may be indicated following failure of lithotripsy or if the size of an intraglandular stone reaches 12 mm or more as the success of

lithotripsy may be less than 20% in such cases⁴. Parotid gland removal should only be carried out for cases of sialolithiasis resistant to minimally invasive techniques.

References

1. Leung A K, Choi M C, Wagner G A. Multiple sialoliths and a sialolith of unusual size in the submandibular duct. *Oral Surg, Oral Med, Oral Path, Oral Radiol, Endo* 1999;87:331-333.
2. Cawson R A, Odell E W. *Essentials of oral pathology and oral medicine*. 6th ed. pp 239-240. Edinburgh: Churchill Livingstone, 1998.
3. Steiner M, Gould A R, Kushner G M, Weber R, Pesto A. Sialolithiasis of the submandibular gland in an 8-year-old child. *Oral Surg, Oral Med, Oral Pathol, Oral Radiol, Endod* 1997;83:188.
4. Zenk J, Benzl W, Iro H. New modalities in the management of human sialolithiasis. *Minimally invasive therapy* 1994;3:275-284.
5. Tejan M G, Rohiwal R L. Multiple salivary calculi in Wharton's duct. *J Laryngol Otol* 1985;99:1313-1314.
6. Williams M F. Sialolithiasis. *Otolaryngologic Clinics of North America* 1999;32:819-834.
7. Akin I, Esmer N. A submandibular sialolith of unusual size: a case report. *J Otolaryngol* 1991;20:123-125.
8. Hoggins G S. Large calcified mass in the sub maxillary gland. *Oral Surg, Oral Med, Oral Pathol* 1968;25:679-681.
9. Naraynsingh V. Giant submandibular gland calculi. *J Oral Maxillofac Surg* 1985;43:384-385.
10. Zakaria M A. Giant calculi of the submandibular salivary gland. *Br J Oral Surg* 1981;19:230-232.
11. Raskin S Z, Gould S M, Williams A C. Submandibular duct sialolith of unusual shape and size. *J Oral Surg* 1975;33:142-145.
12. Mustard T A. Calculus of unusual size in Wharton's duct. *Br Dent J* 1945;79:129.
13. Rust T A, Messerly C D. Oddities of salivary calculi. *Oral Surg, Oral Med, Oral Pathol* 1969;28:862-865.
14. Brusati R, Fiamminghi L. Large calculus of the submandibular gland: report of case. *J Oral Surg* 1973;31:710-711.
15. Carr S J. Sialolith of unusual size and configuration. *Oral Surg, Oral Med, Oral Pathol* 1965;20:709-712.
16. Marchul F, Kurt A M, Dulguerov P, Lehmann W. Retrograde Theory in sialolithiasis formation. *Archives of Otolaryngology- Head and Neck Surgery* 2001;127:66-8.
17. Rauch S, Gorlin R J. Disease of the salivary glands. In: Gorlin R J, Goldmann H M, eds. *Thomas' Oral Pathology*. St Louis, Mo: Mosby-Year Book Inc; 1970:997-1003.
18. Pollack C V Jr, Severance H W Jr. Sialolithiasis: case studies and review. *J Emergency Medicine* 1990;8:561-565.
19. Pietz D M, Bach D E. Submandibular sialolithiasis. *General Dentistry* 1987;35:494-496.
20. Rice D H. Advances in diagnosis and management of salivary gland diseases. *West J Med* 1984;140:238-249.
21. Blatt I. Studies in sialolithiasis: III. Pathogenesis, diagnosis, and treatment. *South Med J* 1964;57:723-728.
22. Zenk J, Werner G, Hosemann M D, Iro H. Diameters of the main excretory ducts of the adult human submandibular and parotid gland – a histological study. *Oral Surg, Oral Med, Oral Pathol, Oral Radiol, Endod* 1998;85:576-580.

23. Work W P, Hecht D W. Inflammatory diseases of the major salivary glands. In Papperalla M M, Shumrick DF (eds): Otolaryngology, Vol 3, 1980. Philadelphia, W B Saunders, 2235-2243.
24. Sherman J A, Mc Gurk M. Lack of correlation between water hardness and salivary calculi in England. *Br J Oral Maxillofac Surg* 2000;38:50-53.
25. Lustmann J, Regev E, Melamed Y. Sialolithiasis: A survey on 245 patients and a review of the literature. *Int J Oral Maxillofac Surg* 1990;19:135-138.
26. Ho V, Currie W J R, Walker A. Sialolithiasis of minor salivary glands. *Br J Oral Maxillofac Surg* 1992;30:273-275.
27. Isacson G, Isberg A, Haverling M, et al. Salivary calculi and chronic sialadenitis of the submandibular gland: A radiographic and histological study. *Oral Surg* 1984;58:622-627.
28. Mc Gurk M, Esudier M. Removing salivary stones. *Br J Hosp Med* 1995;54:184-185.
29. Iro H, Schneider H Th, Fodra C et al. Shockwave lithotripsy of salivary duct stones. *Lancet* 1992;339:1333-1336.
30. Nahlieli O, Baruchin A M. Long-term experience with endoscopic diagnosis and treatment of salivary gland inflammatory diseases. *Laryngoscope* 2000;110:988-993.
31. Nahlieli O, Baruchin A M. Sialendoscopy: three years' experience as a diagnostic and treatment modality. *J Oral Maxillofac Surg* 1997;55:912-918.