

Mandibular cortical index as a predictor of skeletal osteoporosis - a panoramic radiographic study

Suma MS¹, Sujatha GP², Ashok L^{3,*}, Akhilanand Chaurasia⁴

¹Reader, Subbaiah Institute of Dental Sciences, Karnataka, ²Professor, ³Professor & HOD, Bapuji Dental College & Hospital, ⁴Assistant Professor, Dept. of Oral Medicine & Radiology, Faculty of Dental Sciences, King George's Medical University, Lucknow

***Corresponding Author:**

Email: ashok_l2002@yahoo.com

Abstract

Objectives: The objective of our study was to measure Radiomorphometric Indices of mandible on hundred Dental Panoramic Radiographs (DPR) taking into account, age and gender of participants and comparing with Ultrasonometry of heel bone mineral density (Calscan) measurements. Also to establish a precise technique for measuring bone mineral density (BMD) in elderly using DPR so as to recommend DPR as a simple screening aid in osteoporotic risk individual in order to prevent and for better management of osteoporosis.

Methods: The study group consisted of hundred subjects with sixty female and forty male subjects who were apparently healthy without any systemic medication. BMD of the heel was assessed by Quantitative ultrasound. Mandibular Cortical Index [MCI] was calculated from measurements obtained from DPR by two examiners. Both inter and intra examiner's reliability & results of variables such as age and sex were statistically analysed and correlated with Calscan readings.

Results: Results of our study indicated age related bone loss as indicated by calscan. Chi-square analysis for the correlation of MCI with Calscan was found to be 12.41 with significant P value. MCI had sensitivity of 88%, specificity of 39%, positive predictive value of 91%, and negative predictive value of 33% with an overall efficiency of 82% in detecting osteoporosis. But there was no significant correlation found between MCI with age and sex of subjects.

Conclusion: It can be concluded that MCI aids in early diagnosis of osteoporotic patients in the dental office as being at risk for osteoporosis.

Keywords: DPR, MCI, Calscan, BMD.

Introduction

The loss of bone mass in human beings with increasing age is a universally observed phenomenon.^{1,2} Osteopenia and Osteoporosis are two terminologies to describe the decrease in bone mass. Osteopenia can be identified radiographically by a reduction in radiopacity of bone and by observation of thinned cortices, porosity of cortices, or changes in trabecular pattern on panoramic radiographs.² Osteoporosis is a disease of bone that leads to an increased risk of fracture.³ Radiographic assessment of 'bone quality' has applications in implantology and in research, assessing the relationship between oral bone loss and osteoporosis. This aids in early prevention and better management of osteoporosis.⁴

Access to screening for osteopenia or osteoporosis is often limited and the dentist is often the most regularly visited doctor in the elderly population and DPRs are the most frequently used imaging modalities for these patients.⁵ Bone mass has been measured by several techniques.³ DXA including central and peripheral devices, offers the best means of obtaining information on bone mineral density.^{6,7} Quantitative Ultrasound (QUS) has recently been used to assess skeletal status in osteoporosis. QUS has the advantage in that they do not involve ionizing radiation, economical, safe^{8,9} and may provide information on the structural organization of bone in addition to bone

mass.^{7,10} Thus QUS serves as an excellent aid for screening osteoporosis.⁷ QUS techniques have been evaluated in a large number of studies. Their use has been best established for calcaneal systems.⁶ Its low cost and portability make QUS more attractive for use in assessing the risk of fractures in larger populations than may be appropriate for bone DXA.⁹

Various studies have demonstrated that osteoporotic individuals have altered morphology of the mandible and also there exists a good correlation between mandibular and skeletal bone mineral densities.¹¹ Dental radiographs, and especially panoramic images, have been used to predict patients with low BMD.^{12,13} By using various indices, it is possible to measure bone mineral density of mandible.³ Most studies have focused on thickness and integrity of the inferior border of the mandible.

Hence our study is intended to know the efficacy of Dental Panoramic radiographs in predicting osteoporotic risk individuals as compared with bone mineral density of the heel (calscan).

Methodology

This study was conducted on outpatients in the Department of Oral Medicine and Radiology. Hundred normal individuals with female patient's age ranging from forty five to sixty five years and male patient's age ranging from fifty to sixty five years who fulfilled

the criteria of our study were selected. Taking the consent of the patient, relevant information regarding habits, family history, medical history and drug history; the gender and age of the patient were recorded before subjecting them to panoramic radiographs and bone densitometry. Depending on the age, sex and built of the patient, the exposure parameters were kept between 70kv-80kv, current of 10 mA with exposure time of 12 sec.

Measurements of parameters

All processed panoramic radiographs were viewed for inferior cortical border of mandible. Radiographs were traced for inferior cortical border of mandible (Photograph 1).

Two sets of measurements were made by two examiners, 'A' and 'B'.


Measurements of mandibular cortical index (MCI)

Classification of MCI (Klemetti's index) was based on the changes in the inferior cortex on panoramic radiographs as C1, C2, and C3 by using Klemetti's classification (Photograph 1) as follows.^{3,4}

C1- the endosteal margin of the inferior cortex is smooth on both ends.

C2 - the endosteal margin shows semilunar defects or appears to form endosteal cortical residues.

C3- the cortex is obviously porous with dense endosteal residues.



Photograph 1: Reveals a traced panoramic radiograph with endosteal residues in the lower cortical border of mandible

Calscan (Ultrasonometry of Heel)

In order to measure the skeletal BMD, Calscan was done by Densitometer CM100, a peripheral ultrasound bone densitometer with electric power 100vac to 60 VA, mean frequency of 500 kHz and measuring time approximately 10 sec. The attenuation of this sound was recorded by the densitometer which gives printed T score reading (Photograph 2). The name, age and sex of the subjects were filled in the proforma and entered in Microsoft EXCEL spread sheet which was further

subjected to statistical analysis using one way ANOVA, students "t" test, Pearson's correlation and Kappa measure of agreement.



Photograph 2: Reveals a female patient with calscan procedure

Results

According to Calscan, among the study subjects, 13% subjects had normal BMD, 38% subjects were osteopenic and 49% subjects were osteoporotic [Pie diagram 1; Table 1]. The chi-square analysis (X^2) was carried out to find correlation between Calscan and age and was found to be 13.06 and $P < 0.05$ [Tables 2]. The chi-square analysis (X^2) was also carried out to find correlation between Calscan and sex, which was found to be 9.19 and $P < 0.05$. There was a significant correlation observed between Calscan measurements with both age and sex [Tables 3]. MCI had found 15% subjects as C1 (normal), 18% subjects as C2 (osteopenic) and 67% subjects as C3 (osteoporotic) [Table 1]. The chi-square analysis (X^2) was carried out to find correlation between MCI and age and was found to be 5.69 and $P = 0.46$. Hence no positive correlation found between MCI and age of the subjects [Tables 2; Bar diagram 1]. The chi-square analysis (X^2) was carried out to find correlation between MCI and sex and was found to be 3.33 and $P = 0.19$. There was no positive correlation found between MCI and sex of the subjects [Tables 3; Bar diagram 2].

Table 1: Classification of subjects based on Calscan findings

Calscan findings	T score	Number of patients	Range
Normal	> - 1	13 (13%)	2.1 to -0.9
Osteopenia	-1 to -	38 (38%)	-1.1 to -2.4
Osteoporosis	< - 2.5	49 (49%)	-2.5 to -5.0
Total		100	

Table 2: Calscan and MCI in relation to Age

Age (years)	Number	Calscan			MCI		
		Normal	Osteopenia	Osteoporosis	C1	C2	C3
45-49	20	2 (10%)	12 (60%)	6 (30%)	4 (20%)	3 (15.0%)	13 (65.0%)
50-54	29	6 (20.7%)	8 (27.6%)	15 (51.7%)	7 (24.1%)	3 (10.3%)	19 (65.5%)
55-59	22	4 (18.2%)	10 (45.5%)	8 (36.4%)	2 (9.1%)	5 (22.7%)	15 (68.2%)
60-65	29	1 (3.4%)	8 (27.6%)	20 (69%)	2 (6.5%)	7 (4.1%)	20 (69.0%)
Total	100	13 (13.0%)	38 (38.0%)	49 (49.0%)	15 (15%)	18 (18%)	67 (67.0%)
$X^2 = 13.06$ P < 0.05, significant					$X^2 = 5.69$ P = 0.46, not significant		

Table 3: Calscan and MCI in relation to Sex


Sex	Number	Calscan			MCI		
		Normal	Osteopenia	Osteoporosis	C1	C2	C3
Male	40	10 (25%)	15 (37.5%)	15 (37.5%)	4 (10%)	5 (12.5%)	31 (77.5%)
Female	60	3 (5%)	23 (38.3%)	34 (56.7%)	11 (18.3%)	13 (21.7%)	36 (60%)
Total	100	13	38	49	15	18	67
$X^2 = 9.19$ P < 0.05, significant					$X^2 = 3.33$ P = 0.19, not significant		




Table 4: MCI in relation to Calscan findings

Calscan findings	Number	MCI		
		C1	C2	C3
Normal	13	5(38.5%)	2(15.4%)	6(46.2%)
Osteopenia	38	1(2.6%)	10(26.3%)	27(71.1%)
Osteoporosis	49	9(18.4%)	6(12.2%)	34(69.4%)
Total	100	15	18	67
$X^2 = 12.41$ P < 0.05 Significant				

Table 5: Diagnostic efficiency of MCI for Osteoporosis

MCI	CALSCAN		
	Osteoporosis	Normal	
C2+C3	77	8	85
C1	10	5	15
Total	87	13	100

Pie Diagram 1: Calscan findings

Bar Diagram 1: MCI findings in relation to Age**Bar Diagram 2: MCI findings in relation to Sex****Bar Diagram 3: Calscan & MCI findings****MCI in relation to calscan findings**

MCI had detected 38.5% of subjects in normal group, 2.6% of subjects in osteopenic group and 18.4% of subjects in osteoporotic group as C1. It had also found 15.4% of subjects in normal group, 26.3% of subjects in osteopenic group and 12.2% of subjects in osteoporotic group as C2. MCI had detected 46.2% of subjects in normal group, 71.1% of subjects in osteopenic group and 69.4% of subjects in osteoporotic group as C3. Chi-square analysis was found to be 12.41

whereas $P < 0.05$. There was a highly significant correlation found between MCI and Calscan [Table 4; Bar diagram 3].

Diagnostic efficiency of MCI for osteoporosis

Out of 100 subjects, MCI had detected 85 subjects as diseased (C2+C3) that is osteoporotic and 15 subjects as normal (C1). Over all Calscan had detected 87 subjects as osteoporotic and 13 subjects as normal subjects. The sensitivity of MCI in detecting osteoporosis was 88% whereas it had specificity of 39% with positive predictive value of 91%, negative predictive value of 33% and overall efficiency of 82% [Table 5].

Discussion

In our study, it was observed that with increasing age (69%) of the subjects and also in female subjects (56.7%); there was increase in occurrence of osteoporosis. However the disease was less common in males (37.5%), and also in younger age group (30%). This was comparable to the reports of various studies that are available in the literature.^{14,15,16,17} Our results were in accordance with the results of Nan-Ping Yang et al.¹⁸

Also in our study, there was an increase in the MCI (C3) as age advances and also in male subjects (77.5%), it was more prevalent as compared to females (60%) which was in contrary to other reports. There was positive correlation ($X^2 = 12.41$) between Calscan and MCI with significant P value. The sensitivity of MCI in our study in detecting osteoporosis was 88% where as it had specificity of 39% with positive predictive value of 91%, negative predictive value of 33% and overall efficiency of 82%. Results of our study were comparable to studies^{5,12,19,20,21} and were in contradiction to studies reported in literature.^{3,4,22,23,24}

As observed in our study, there was more number of C1 subjects in normal group than in osteoporotic group and more number of C2 subjects in normal group than in osteoporotic group. Interestingly there was less number of C3 subjects in normal group than in osteoporotic group. Calscan was able to detect the presence of disease in the study group. There was very good correlation found between MCI with Calscan with highly significant P value.

Our results were in accordance with the results of Klemetti E²⁵ who reported that reproducibility of mandibular cortical index was 98% and observed a significant correlation between their classification and vertebral bone mineral density (BMD) as assessed dual energy xray absorptiometry (DXA). Similar results were observed in our study also. Author also reported that sensitivity and specificity of the combination of mandibular cortical width and their classification of mandibular cortical morphology (MCI) in the identification of osteoporosis was 0.13 and 0.99. Similar findings were observed in our study also where

there was a significant correlation found between MCI and Calscan.

A study was conducted in which short-term precision and long-term precision of ultrasound parameters were studied and also relationships between ultrasound parameters and BMD at various sites were examined in 42 subjects. BMD at the heel was assessed at the location corresponding to that of the ultrasound measurements (BMD heel/calscan). It was concluded that ultrasound measurements at the heel correlate well to BMD at the same site.²⁶ Hence we too used calscan for assessing BMD of heel, as it is easily accessible and correlates well with the central bone mineral density (femoral neck).

Conclusion

Results of our study indicated age related bone loss (osteoporosis) as MCI was showing significant correlation with Calscan. It was also observed that the MCI was not correlating with age and sex of the subjects. But there was positive trend of values towards osteoporosis. This is suggesting the need for larger sample size and the more precise technique of measuring skeletal mineral density by using central DXA.

Careful examination of the lower cortical border of mandible in panoramic radiographs helps to classify subjects based on Klemetti's classification as C1, C2 and C3. Wherein C3 being osteoporotic, needs treatment at the earliest there by making individual less susceptible to fractures. Hence Mandibular Cortical Index (MCI) aids in early diagnosis of osteoporotic patients in the dental office as being at risk of osteoporosis, so that they can be referred to a physician for appropriate evaluation.

Acknowledgement

My heartfelt gratitude to all the faculty of department of oral medicine & diagnostic radiology of Bapuji Dental College & Hospital, for helping me in conducting this study.

References

1. Benson B. W, Prihoda, T. J. and Glass B. J. Variations in adult cortical bone mass as measured by a panoramic mandibular index. *Oral Surgery, Oral Medicine, Oral Pathology.*1991;71:349-356.
2. Ivona Musa, Dubravka Knezovic-Zlataric, Asja Celebic, Andrija Bosnjak. The influence of gender and age on the values of linear radiomorphometric indices measured on the lower border of the mandible. *Acta Stomat Croat.*2002;36(2):199-202.
3. Taguchi A, Sueti Y, Ohtsuka M, Otani K, Tanimoto K, Ohtaki M. Usefulness of panoramic radiography in the diagnosis of postmenopausal osteoporosis in women. Width and morphology of inferior cortex of the mandible. *Dentomaxillofac Radiol.*1996;25:263-267.
4. Horner K and Devlin H. The relationships between two indices of mandibular bone quality and bone mineral density measured by dual energy X-ray Absorptiometry. *Dentomaxillofacial Radiology.*1998;27:17-21.
5. Anders Halling, G. Rutger Persson, Johan Berglund, Owe Johansson, Stefan Renvert. Comparison between the Klemetti index and heel DXA BMD measurements in the diagnosis of reduced skeletal bone mineral density in the elderly. *Osteoporosis International.*2005;16(4):999-1003.
6. He YQ, Fan B, Hans D, Li J, Wu CY, Njeh CF et al. Assessment of a new quantitative ultrasound calcaneus measurement: precision and discrimination of hip fractures in elderly women compared with dual X-ray absorptiometry. *Osteoporos Int.* 2000;11(4):354-60.
7. Sheryl L. Follin, Laura B. Hansen. Current approaches to the prevention and treatment of postmenopausal osteoporosis. *American Journal of Health-System Pharmacy.*2003;60(9):883-901.
8. S Cheng, C F Njeh, B Fan, X Cheng, D Hans, L Wang et al. Influence of region of interest and bone size on calcaneal BMD: implications for the accuracy of quantitative ultrasound assessments at the calcaneus. *The British Journal of Radiology.*2002;75:59-68.
9. Maghraoui A, Morjane F, Mounach A, Ghazi M, Nouijai A, Achemlal L. et al. Performance of calcaneus quantitative ultrasound and dual-energy x-ray absorptiometry in the discrimination of prevalent asymptomatic osteoporotic fractures in postmenopausal women. *Rheumatology International.* 2009;29(5):551-556.
10. Karen M Knapp. Quantitative ultrasound and bone health. *Salud Publica de Mexico.* 2009;51(1):S18-S24.
11. Mohajery M, Brooks S L. Oral radiographs in the detection of early signs of osteoporosis. *Oral Surg Oral Med Oral Pathol.*1992;73:112-7.
12. B Cakur, S Dagistan, A Sahin, O Altun, F Caglayan, O Miloglu et al. Dental Panoramic Radiography in the Diagnosis of Osteoporosis. *The Journal of International Medical Research.*2008;36:792-799.
13. Horner K, Devlin H, Alsop CW, Hodgkinson IM, Adams JE. Mandibular bone mineral density as a predictor of skeletal osteoporosis. *British Journal of Radiology.*1996;69(827):1019-1025.
14. Stefano Gonnelli, Chiara Cepollaro, Luigi Gennari, Andrea Montagnani, Carla Caffarelli, Daniela Merlotti et al. Quantitative ultrasound and dual-energy X-ray absorptiometry in the prediction of fragility fracture in men. *Osteoporos Int.* 2005;16:963-968.
15. Ishikawa K, Ohta T. Radial and metacarpal bone mineral density and calcaneal quantitative ultrasound bone mass in normal Japanese women. *Calcif Tissue Int* 1999;65:112-116.
16. Von Wower N. Bone mineral content of mandibles: normal reference values—rate of age-related bone loss. *Calcif Tissue Int.* 1988;43:193-198.
17. Chan WP, Liu JF, Chi WL. Evaluation of bone mineral density of the lumbar spine and proximal femur in population-based routine health examinations of healthy Asians. *Acta Radiol.* 2004;45:59-64.
18. Nan-Ping Yang, Ian Jen, Shao-Yuan Chuang, Shui-Hu Chen, and Pesus Chou. Screening for low bone mass with quantitative ultrasonography in a community without dual-energy X-ray absorptiometry: population-based survey. *Musculoskelet Disord.* 2006;7-24.
19. Vinicus Dutra, Hugh Devlin, Cristiano Sosin, Jie Yang, Keith Horner and Arthur Rocha Corea Fernandes. Mandibular morphological changes in low bone mass edentulous females; evaluation of panoramic radiographs. *Oral Surg Oral Med Oral Pathol Oral Endo.*2006;102:663-8.

20. Zlataric DK, Celebic A. Clinical bone densitometric evaluation of the mandible in removable denture wearers dependent on the morphology of the mandibular cortex. *J Prosthet Dent.*2003;90:86–91.
21. Knezovic Zlataric D, Celebic A, Lazic B, Baucic I, Komar D, Stipetic-Ovcaricek J et al. Influence of age and gender on radiomorphometric indices of the mandible in removable denture wearers. *Coll Antropol.* 2002;26:259–266.
22. B Cakur, S Dagistan, A Sahin, A Harorli and A B Yilmaz. Reliability of mandibular cortical index and mandibular bone mineral density in the detection of osteoporotic women. *Dentomaxillofacial Radiology.*2009;38:255–261.
23. Diez-Perez A, Marin F, Vila J, Abizanda M, Cervera A, Carbonell C et al. Evaluation of calcaneal quantitative ultrasound in a primary care setting as a screening tool for osteoporosis in postmenopausal women. *J Clin Densitom.* 2003 Fall;6(3):237-45.
24. Akira Taguchi, Mikio Tsuda, Masahiko Ohtsuka, Ichiro Kodama, Mitsuhiro Sanada, Takashi Nakamoto et al. Use of dental panoramic radiographs in identifying younger postmenopausal women with osteoporosis. *Osteoporos Int.* 2006;17:387–394.
25. Graafmans WC, Van Lingen A, Ooms ME, Bezemer PD, Lips P. Ultrasound measurements in the calcaneus: precision and its relation with bone mineral density of the heel, hip, and lumbar spine. *Bone.* 1996 Aug;19(2):97-100.
26. Klemetti E, Kolmakov S, Kroger H. Pantomography in assessment of the osteoporosis risk group. *Scand J Dent Res.* 1994:68–72.